

## ***Pharmacy Provider Reminder, ICD9 Code Requirement***

***This is a reminder notice to the MS Division of Medicaid (DOM) pharmacy providers regarding ICD9 requirements at point of sale or POS. Please note that the article in its entirety was published in the June 2012 Provider Bulletin.***

***Be advised that claims for the drugs below will deny without the validating ICD9 codes. This includes Suboxone/Subutex. Refer to directions below regarding claim submission.***

### ***ICD-9 Codes at Pharmacy Point of Sale (POS)***

Effective July 1, 2012, the Division of Medicaid will accept specific ICD-9 codes in the pharmacy point of sale (POS) system for the following drugs ***only***:

- Xarelto –limited to ***hip replacement; and knee replacement***
- Effient –limited to ***unstable angina; STEMI*** (Segment Elevation Myocardial Infarction or heart attack), ***NSTEMI*** (Non-ST Segment Elevation Myocardial Infarction)
- Brilintia—limited to ***unstable angina; STEMI*** (Segment Elevation Myocardial Infarction or heart attack), and ***NSTEMI*** (Non-ST Segment Elevation Myocardial Infarction)
- Suboxone/Subutex—limited to ***opioid dependence***

Prescriber must write the validating ICD code on the prescription and no manual prior authorization is required. For the comprehensive list of valid ICD codes at POS, refer to Pharmacy Services' webpage at <http://www.medicaid.ms.gov/Pharmacy.aspx>, and select POS ICD-9 Codes. For other indications for the aforementioned drugs, please submit a prior authorization request.

To process a claim, the pharmacy is to use the following procedure:

- (1) Submit the diagnosis code (**on RX**) in the field **Diagnosis Code (492-DO)**
- (2) Submit qualifier **Diagnosis Code Qualifier (492-WE) = 01**

Field	Field Name	Values Supported
492- WE	Diagnosis Code Qualifier	Required when Diagnosis used. <b>01</b> = ICD9
492-DO	Diagnosis Code	Required when diagnosis is needed for designated drug coverage.